

**CHEM 3311**

**HARRINGTON**

**Exam 1 7:00 – 8:30 PM February 14, 2017 in HUMN1B50**

**Instructions.** No notes, books, laptops, phones, calculators, models or drawing stencils.

Periodic Table and electronegativity chart are provided.

**NAME:**

	<b>Points Possible</b>	<b>Score</b>
<b>1</b>	12	
<b>2</b>	15	
<b>3</b>	16	
<b>4</b>	16	
<b>5</b>	12	
<b>6</b>	16	
<b>7</b>	13	
<b>Exam 1 Total Raw Score</b>	100	
<b>Curve</b>		
<b>Exam 1 Curved Score</b>		
<b>Exam 1 Letter Grade</b>		

1. (12 points) Complete the Lewis Dot structure for each of the molecules below. Calculate the formal charge on each atom in the completed Lewis Dot structure.

**NOTE:** For molecules which can be represented by resonance hybrids, draw only the most important contributor to the hybrid.

	<b>Formula</b>	<b>Lewis Dot Structure</b>
--	----------------	----------------------------




a.	CS <sub>2</sub>	
----	-----------------	--

b.	HN <sub>3</sub>	
----	-----------------	--

c.	O <sub>3</sub>	
----	----------------	--

d.	C <sub>2</sub> F <sub>4</sub>	
----	-------------------------------	--


2. (15 points) For each resonance hybrid provided, draw the two-electron curved arrows that show the conversion of the first contributor to the second. If there is a third contributor, also draw the two-electron curved arrows that show the conversion of the second to the third. Identify the most important contributor to the hybrid and provide a complete evaluation of each contributor to explain why it is the most important.



3. (16 points) Draw a structure for 3-methylpentane.

Complete the Newman Projections for the conformations of 3-methylpentane obtained by rotation about the C2-C3 bond.

(Hint: Rotate the back C 60° clockwise to convert the projection A to B, B to C, etc.)



Use the eclipsing and gauche strain data in the **Tables** below to determine which conformation has the least strain and which conformation has the most strain.

**Table 1.** Eclipsing Strain Energy (kJ/mol)

**Table 2.** Gauche Strain Energy (kJ/mol)

	<b>H</b>	<b>CH<sub>3</sub></b>	<b>CH<sub>3</sub>CH<sub>2</sub></b>			<b>H</b>	<b>CH<sub>3</sub></b>	<b>CH<sub>3</sub>CH<sub>2</sub></b>
<b>H</b>	3.8	5.9	6.3		<b>H</b>	0	0	0.4
<b>CH<sub>3</sub></b>	5.9	10.5	11.3		<b>CH<sub>3</sub></b>	0	3.3	3.8
<b>CH<sub>3</sub>CH<sub>2</sub></b>	6.3	11.3	13.8		<b>CH<sub>3</sub>CH<sub>2</sub></b>	0.4	3.8	4.6

(For example: The strain energy associated with an H eclipsing another H is 3.8 kJ/mol.)

4. (16 points) Draw the structures corresponding to each name.  
Which of the compounds in each group are constitutional isomers?



**Group 1.**      butane                      cyclobutane                      2-methylpropane

**Group 2.**      cyclopentane                      2,2-dimethylpentane                      2,2,3-trimethylbutane

**Group 3.**      cyclohexane                      ethylcyclopentane                      propylcyclopropane


**Group 4.**      1-cyclopropylbutane                      isopropylcyclobutane                      *tert*-butylcyclohexane

5. (12 points) For each group of three compounds, identify the compound with the highest boiling point and the compound with the lowest boiling point. Explain each answer (identify the intermolecular attractive forces for each compound).






7. (13 points) For the reaction:



Consider the reaction coordinate-energy diagram:



Draw the two-electron curved arrows that show the conversion of reactants to products.

Draw the structure of the reactive species present at point **B** on the diagram.

Estimate the activation energy (kJ/mol) for the forward reaction.

Is the forward reaction endothermic or exothermic?

Estimate the ratio of products to reactants at equilibrium.