Easily Legible Printed Name:	
------------------------------	--

CHEM 3371 (100), Spring 2017 Professor Walba First Hour Exam February 14, 2017

scores:

- 1)
- 2)
- 3)
- 4)

5)

,

CU Honor Code Pledge: On my honor, as a University of
Colorado at Boulder Student, I have neither given nor received
unauthorized assistance.


Signature:

Recitation TA Name: _____ Ed Guzman

Recitation day and time:

This is a closed-book exam. The use of notes, calculators, scratch paper, or cell phones will not be allowed during the exam. You may use models brought in a clear Ziploc bag. Please put all your answers on the test in the appropriate place. Use the backs of the pages for scratch (there are two additional blank scratch sheets after the last page of the exam). DO NOT PUT ANSWERS ON THE SCRATCH SHEETS.

PLEASE read the questions very carefully!



Page 1 of 7

1 (20 pts) a) For each of the following heterocyclic molecules, indicate whether the molecule is aromatic or not aromatic by circling your answer, give the hybridization of the heteroatom in the box, and give a **brief** explanation of your answer in the open space to the right of the structure.


1)	∭"-H
1)	N-H

aromatic

not aromatic



aromatic



not aromatic



3) B-H

aromatic

not aromatic



4) B-H

aromatic

not aromatic



b) Cyclopropenone (compound 1 below) looks extremely strained, but it's actually surprisingly stable (has a surprisingly low heat of formation). Also, the measured dipole moment is larger than one would expect for a ketone. Give a brief explanation of your hypothesis for why this is the case.



1



2) (20 pts) Give the **single major product** for each of the following reactions, carefully showing stereochemistry using wedges and dashes if appropriate. If a racemate is formed, show only one enantiomer and label it "rac." Assume chiral starting materials are single pure enantiomers unless they are labeled "rac."

a)
$$H_2SO_4$$
 (large excess)

b)
$$\frac{\text{cat Hg(OAc)}_2, \text{H}_2\text{O}}{}$$

3) (20 pts) Propose reagents for accomplishing each of the following transformations. For reactions involving sequential addition of reagents, label the two parts of the reaction using letters. Your synthesis may require multiple reactions, with isolation of intermediate products, to make the target. Use numbers to indicate individual steps in a multi-step synthesis. Make your synthesis efficient (i.e. the target product should be the major product). Assume chiral starting materials and products are single pure enantiomers unless they are labeled "rac."

4) (20 pts) Propose an arrow-pushing mechanism for each of the following reactions.

b)
$$H_3C-C\equiv C-CH_3$$
 Na^0 , liquid NH_3 H_3C CH_3

Printed Name:

Q4 – continued

c)
$$H_3C-C\equiv C-H$$
 $cat Hg(OAc)_2$ O H_2SO_4, H_2O

D 1 NT		
Printed Name:		

5) (20 pts) Propose a synthesis of the following diol target using any organic starting materials with **five carbons or less**. You may use any necessary inorganic reagents. Try to make your synthesis efficient (i.e. the target should be produced in high yield). More than one step will be required. Each reaction in the sequence leading to an isolated and purified product should be shown. Please use letters to indicate sequential addition of reagents in a single reaction. **Do not put multiple reactions over/under one arrow!**